

## DAFTAR PUSTAKA

- Alhadi, F., Kaprawi, F., Hamidy, A., & Kirschey, T. (2021). Panduan Bergambar dan Identifikasi Amfibi Pulau Jawa. Perkumpulan Amfibi Reptil Sumatra (ARS/NABU).
- JakartaArifin, U. (2018) ‘Phylogenetic systematics, diversity, and biogeography of the frogs with gastromyzophorous tadpoles (Amphibia: Ranidae ) on Sumatra, Indonesia’, pp. 1–175.
- Arifin, U. (2018). Phylogenetic systematics, diversity, and biogeography of the frogs with gastromyzophorous tadpoles (Amphibia: Ranidae ) on Sumatra, Indonesia. 1–175
- Boulenger, G.A. (1887) ‘On new reptiles and batrachians from North Borneo’, *Annals and Magazine of Natural History, Series 5*, 20, pp. 95–97.
- Che, J. et al. (2007) ‘Phylogeny of Raninae (Anura: Ranidae) inferred from mitochondrial and nuclear sequences’, *Molecular Phylogenetics and Evolution*, 43(1), pp. 1–13. Available at: <https://doi.org/10.1016/j.ympev.2006.11.032>.
- Cisneros-Heredia, D.F. and McDiarmid, R.W. (2007) *Revision of the characters of Centrolenidae (Amphibia: Anura: Athesphatanura), with comments on its taxonomy and the description of new taxa of glassfrogs*, Zootaxa. Available at: <https://doi.org/10.11646/zootaxa.1572.1.1>.
- Dubois, A. (1992) ‘Notes sur la classification des Ranidae (Amphibiens, Anoures)’, *Bulletin mensuel de la Société linnéenne de Lyon*, 61(10), pp. 305–352. Available at: <https://doi.org/10.3406/linly.1992.11011>.
- Firdaus, N. (2013) ‘Meristogenys Yang, 1991 (Anura: Ranidae) from Kalimantan: Conservation, Opportunities, and Threats’, *The 2nd Society for Indonesian Biodiversity International Conference on Biodiversity Volume: Vol 2, July 2013*, 1991(February). Available at: <https://doi.org/10.13140/RG.2.1.2855.8883>.
- Frost, D.R. et al. (2006) ‘The amphibian tree of life’, *Bulletin of the American*

- Museum of Natural History*, 0090(297), pp. 1–370. Available at: [https://doi.org/10.1206/0003-0090\(2006\)297\[0001:TATOL\]2.0.CO;2](https://doi.org/10.1206/0003-0090(2006)297[0001:TATOL]2.0.CO;2).
- Frost, Darrel R. (2024). Amphibian Species of the World: an Online Reference. Version 6.2 (*Date of access*). Electronic Database accessible at <https://amphibiansoftheworld.amnh.org/index.php>. American Museum of Natural History, New York, USA. doi.org/10.5531/db.vz.0001
- Günther, A.C.L.G. (1872) ‘On the reptiles and amphibians of Borneo. Proceedings of the Zoological Society of London 1872’, pp. 586–600. Available at: <http://www.nber.org/papers/w16019>.
- Hall, R. (2013) ‘The palaeogeography of Sundaland and Wallacea since the Late Jurassic’, *Journal of Limnology*, 72(S2), pp. 1–17. Available at: <https://doi.org/10.4081/jlimnol.2013.s2.e1>.
- Hall, R. and Morley, C.K. (2004) ‘Sundaland basins’, *Geophysical Monograph Series*, 149(January 2004), pp. 55–85. Available at: <https://doi.org/10.1029/149GM04>.
- Hamidy, A. and Kurniati, H. (2015) ‘A new species of tree frog genus Rhacophorus from Sumatra, Indonesia (Amphibia, Anura)’, *Zootaxa*, 3947(1), pp. 49–66. Available at: <https://doi.org/10.11646/zootaxa.3947.1.3>.
- Idrus, M.R. *et al.* (2021) ‘Morphological Variations of The Genus Huia Yang, 1991 in Region of Java, Kalimantan and Sumatra’, *Jurnal Biodjati*, 6(1), pp. 1–10. Available at: <https://doi.org/10.15575/biodjati.v6i1.10771>.
- Inger, R. F., and P.A.G. (1983) ‘Variation of Bornean frogs of the Amolops jerboa species group, with description of two new species’, *Zoology*, (Fieldiana), p. New Series 19: 1–13. Available at: <http://www.nber.org/papers/w16019>.
- Inger, R.F. (1966) ‘The systematics and zoogeography of the Amphibia of Borneo’, *Fieldiana. Zoology* 52, pp. 1–402.
- Iskandar, D.T., Bickford, D.P. and Arifin, U. (2011) ‘A new Ingerana (Anura, Dicroglossidae) with no external tympanum from Borneo, Indonesia’, *Raffles Bulletin of Zoology*, 59(2), pp. 213–218.
- Kurniawan, N. et al. (2011) ‘Taxonomic status of three types of Fejervarya

- cancrivora from Indonesia and other Asian countries based on morphological observations and crossing experiments’, *Zoological Science*, 28(1), pp. 12–24. Available at:<https://doi.org/10.2108/zsj.28.12>.
- Kurniawan Nia, Ahmadlia Driyana Rike, Nahari Day Shine, Firdaus Anggun Sausan. (2015) ‘Speciation And Zoogeography Of Amphibian In Sundaland’, 21(1), pp.1–7.
- Kuspriyanto (2015) ‘Upaya konservasi keanekaragaman hayati dikawasan lindung di Indonesia’, *Metafora*, 1(2), pp. 134–142.
- Liu, C.C. (1950) ‘Amphibians of western China. Fieldiana: Zoology’, *Memoir*, 2: 1-400.
- Lohman, D.J. et al. (2011) ‘Beyond Wallaces line: Genes and biology inform historical biogeographical insights in the Indo-Australian archipelago’, *Annual Review of Ecology, Evolution, and Systematics*, 42(May 2014). Available at: <https://doi.org/10.1146/annurev-ecolsys-102710-145001>.
- Malkmus, R. et al. (2002) *Amphibians & reptiles of Mount Kinabalu (North Borneo)*, *Amphibians & reptiles of Mount Kinabalu*.
- Matsui, M. (1984) ‘Morphometric variation analyses and revision of the Japanese toads (genus *Bufo*, *Bufonidae*)’, *Contributions from the Biological Laboratory, Kyoto University*, 26, pp. 209–428. Available at: <http://repository.kulib.kyoto-u.ac.jp/dspace/handle/2433/156031>.
- Matsui, M. (1986) ‘Three New Species of Amolops from Borneo (Amphibia , Anura , Ranidae) Author ( s ): Masafumi Matsui Published by : American Society of Ichthyologists and Herpetologists ( ASIH ) Stable URL : <https://www.jstor.org/stable/1444943> REFERENCES Linked referen’, 1986(3), pp. 623–630.
- Matsui, M. et al. (2006) ‘Phylogenetic relationships of Oriental torrent frogs in the genus Amolops and its allies (Amphibia, Anura, Ranidae)’, *Molecular Phylogenetics and Evolution*, 38(3), pp. 659–666. Available at: <https://doi.org/10.1016/j.ympev.2005.11.019>.
- Matsui, M. et al. (2024) ‘Revision of the *Limnonectes kuhlii*-like Fanged Frogs

from Malaysian Borneo (Amphibia: Anura: Dic平glossidae)', *Current Herpetology*, 43(2), pp. 226–277. Available at: <https://doi.org/10.5358/hsj.43.226>.

Masafumi Matsui, Tomohiko Shimada, and Ahmad Sudin "A New Species of Meristogenys (Amphibia, Anura, Ranidae) from Sabah, Borneo," *Zoological Science* 27(1), 61-66, (1 January 2010).

McLeod, D.S. et al. (2011) "Same-same, but different": An unusual new species of the *Limnonectes kuhlii* Complex from West Sumatra (Anura: Dic平glossidae)', *Zootaxa*, (2883), pp. 52–64. Available at: <https://doi.org/10.11646/zootaxa.2883.1.4>.

Mocquard F. (1890). Recherches sur la faune herpetologique des Iles de Bornéo et de Palawan. Nouvelles Archives du Muséum d'Histoire Naturelle, Série 3 2: 115–168, pls 7–11.

Prafiadi, S., Kurniawan, N. and Hamidy, A. (2016) 'Keberagaman Spesies Katak Pohon Hijau Papua *Litoria infrafrenata* *infrafrenata* Tyler, 1971 pada Wilayah Kepulauan Wallacea dan Indo- Australia', *Jurnal Pembangunan dan Alam Lestari (J-PAL)*, 7(1), pp. 33–43. Available at: <https://jpdl.ub.ac.id/index.php/jpal/article/view/214>.

Sanders, M. (2021) *Photographic Field Guide to Australian Frogs, Photographic Field Guide to Australian Frogs*. Available at: <https://doi.org/10.1071/9781486313266>.

Sarwenda, Subagio and Imran, A. (2016) 'Struktur Komunitas Amphibi Di Taman Wisata Alam (TWA) Kerandangan Dalam Upaya Penyusunan Modul Ekologi Hewan', *Jurnal Ilmiah Biologi*, 4(1), pp. 21–26.

Savage, J.M. and Heyer, W.R. (1967) 'Beitrag zur Neotropischen Fauna Variation and distribution in the tree - frog genus *Phyllomedusa* in Costa Rica , central America Variation and distribution in the tree-frog genus *Phyllomedusa*', *Beitrage zur Neotropischen Fauna*, 5(2), pp. 111–131.

Setiawan, A. (2022) 'Keanekaragaman Hayati Indonesia: Masalah dan Upaya Konservasinya', *Indonesian Journal of Conservation*, 11(1), pp. 13–21.

Available at: <https://doi.org/10.15294/ijc.v11i1.34532>.

Shimada, T., (2007) ‘Identity of larval Meristogenys from a single stream in Sabah, Malaysia (Amphibia: Ranidae)’, *Zoological Journal of the Linnean Society*, 151(1), pp. 173–189. Available at: <https://doi.org/10.1111/j.1096-3642.2007.00319.x>.

Shimada, T. et al. (2011) ‘A survey of morphological variation in adult *meristogenys amoropalamus* (Amphibia, Anura, Ranidae), with a description of a new cryptic species’, *Zootaxa*, 56(2905), pp. 33–36. Available at: <https://doi.org/10.11646/zootaxa.2905.1.3>.

Shimada, T. et al. (2015) ‘A new species of meristogenys (Anura: Ranidae) from Sarawak, Borneo’, *Zoological Science*, 32(5), pp. 474–484. Available at: <https://doi.org/10.2108/zs140289>.

Solihuddin, T. (2014) ‘A Drowning Sunda Shelf model during Last Glacial Maximum (LGM) and Holocene: A review’, *Indonesian Journal on Geoscience*, 1(2), pp. 99–107. Available at: <https://doi.org/10.17014/ijog.v1i2.182>.

Surján, G. (2008) ‘Dichotomy - A forgotten ancient principle’, *Studies in Health Technology and Informatics*, 136(June), pp. 869–874. Available at: <https://doi.org/10.3233/978-1-58603-864-9-869>.

Yang, D. -t. (1991) ‘Phylogenetic systematics of the Amolops group of ranid frogs of southeastern Asia and the Greater Sunda Islands.’, *Zoology*, p. New Series 63: 1–42.

